

PRINT your name and student ID: _____

4. Splotchy Writing (10 points)

Professor Courtade writes with a sharpie to accommodate the vision of as many people as possible. Unfortunately, some characters get smudged, which makes them difficult to read. The following is a (hypothetical) passage from lecture notes, and the smudges are labeled (1), (2), ..., (10). Your task is to identify correct expressions for each of the smudges.

Let $A \in \mathbb{R}^{n \times (1)}$ be a matrix with rank r . It is always possible to write A in terms of its compact SVD

$$A = U \Sigma V^T,$$

where Σ is a diagonal $r \times r$ matrix, and $U \in \mathbb{R}^{(2) \times (3)}$ and $V \in \mathbb{R}^{(4) \times (5)}$ have orthonormal columns. This means that $U^T U = I_{(6)}$ and $V^T V = I_{(7)}$, where we write I_m to denote the $m \times (8)$ identity matrix, for an integer m . The columns of U form a basis for the range of A , which is defined as

$$\text{range}(A) = \{A\vec{x} \mid \vec{x} \in \mathbb{R}^k\}.$$

Note that $\text{range}(A)$ is a subspace of $\mathbb{R}^{(9)}$, which has dimension (10).

Select the values for each smudge from the multiple choice below. For each smudge, completely fill in the circle next to the correct answer. (Hint: Resist the temptation to get distracted by unfamiliar terminology... that isn't what this question is about.)

Concepts: This question tests your understanding of matrix multiplication (specifically, compatibility of dimensions necessary for, and resulting from, matrix-matrix multiplication), as well as dimension of column-space (i.e., range). As suggested by the hint, the technical jargon (such as compact SVD, orthonormal) is completely irrelevant to determining the smudged dimensions.

Answer:

- | | | | | |
|------|--------------------------------------|--------------------------------------|--------------------------------------|--------------------------------------|
| (1) | <input type="radio"/> k | <input type="radio"/> m | <input type="radio"/> n | <input type="radio"/> r |
| (2) | <input type="radio"/> k | <input type="radio"/> m | <input checked="" type="radio"/> n | <input type="radio"/> r |
| (3) | <input type="radio"/> k | <input type="radio"/> m | <input type="radio"/> n | <input type="radio"/> r |
| (4) | <input checked="" type="radio"/> k | <input type="radio"/> m | <input type="radio"/> n | <input type="radio"/> r |
| (5) | <input type="radio"/> k | <input type="radio"/> m | <input type="radio"/> n | <input type="radio"/> r |
| (6) | <input type="radio"/> k | <input type="radio"/> m | <input type="radio"/> n | <input type="radio"/> r |
| (7) | <input type="radio"/> k | <input type="radio"/> m | <input type="radio"/> n | <input type="radio"/> r |
| (8) | <input type="radio"/> k | <input checked="" type="radio"/> m | <input type="radio"/> n | <input type="radio"/> r |
| (9) | <input type="radio"/> k | <input type="radio"/> m | <input type="radio"/> n | <input type="radio"/> r |
| (10) | <input type="radio"/> k | <input type="radio"/> m | <input type="radio"/> n | <input checked="" type="radio"/> r |

PRINT your name and student ID: _____

5. Matrix Inversion (10 points)

You landed your first job at 16Atech (the Bay Area's newest and hottest tech company), and your first assignment is to invert a matrix $A \in \mathbb{R}^{n \times n}$. You say "no problem", and implement Gaussian elimination. You obtain the following reduction of the augmented matrix:

$$[A|I] \longrightarrow [I|P].$$

The dimension n is extremely large, so the computation takes several days to complete, and you give your boss the matrix $P \in \mathbb{R}^{n \times n}$ just minutes before the deadline.

- (a) (2 points) Your boss panics, saying "Oh, no! Your procedure only guarantees that $AP = I$ and not necessarily that $PA = I$." In *one sentence*, concisely explain why your boss thinks this might be an issue.

Concepts: This part tests your understanding of what the reduction $[A|I] \longrightarrow [I|P]$ means in terms of linear equations.

- (b) (8 points) You try to calm them down, saying "Don't worry, the matrix also satisfies $PA = I$, and therefore P is the inverse of A just like you wanted. I'll prove it to you..."

Your proof consists of the following two steps (fill in the details as your answer to this question):

Step 1: Argue that your matrix P is the unique $Q \in \mathbb{R}^{n \times n}$ satisfying $AQ = I$.

Step 2: Prove that $PA = AP = I$. (Hint: consider the matrix $A(P + PA - I)$)

As suggested by part (a), you should not assume that A^{-1} exists. Proving that it does is the point of this problem.

Concepts: Step 1 asks you to interpret what the augmented matrix $[I|P]$ reveals about number of solutions to the corresponding system of linear equations. Step 2 requires you to use the distributive property of matrix multiplication together with the previously established property of P .

Answer:

Step 1: The reduction $[A|I] \rightarrow [I|P]$ implies there is a unique solution $X = P$ to the system of equations $AX = I$, since there are no free variables.

Step 2: ...

PRINT your name and student ID: _____

6. Tomography (19 points)

Recall that in our simple tomography example of 4 pixels arranged into a 2×2 matrix, our initial set of measurements produced the following system of equations with unknowns x_1, \dots, x_4 and measured intensities b_1, \dots, b_4 :

$$\begin{array}{rcccc} x_1 & +x_2 & & & = b_1 \\ & & x_3 & +x_4 & = b_2 \\ x_1 & & +x_3 & & = b_3 \\ & x_2 & & +x_4 & = b_4 \end{array}$$

- (a) (3 points) Write the above system of equations in matrix-vector form $A\vec{x} = \vec{b}$.

Concepts: Do you know how to write a system of equations in matrix-vector form?

- (b) (8 points) Use Gaussian elimination to find a basis for the nullspace of your matrix in part (a). Show your work.

Concepts: This question tests whether you know the definition of nullspace, and the mechanics of gaussian elimination for solving the system $A\vec{x} = \vec{0}$.

Answer: A basis for $N(A)$ is $[1 \ -1 \ -1 \ 1]^T$.

- (c) (2 points) Suppose \vec{x}_0 denotes the correct pixel values, which of course satisfy $A\vec{x}_0 = \vec{b}$. Give another solution \vec{x}_1 to the system of equations $A\vec{x} = \vec{b}$, satisfying $\vec{x}_1 \neq \vec{x}_0$. Leave your answer in terms of \vec{x}_0 .

Concepts: This question checks whether you understand how nullspace relates to characterizing the set of solutions to a system of linear equations.

Answer: Another solution is $\vec{x}_0 + [1 \ -1 \ -1 \ 1]^T$.

- (d) (2 points) Suppose we add the measurement

$$x_1 + x_4 = b_5.$$

Will the resulting new system of equations always have a solution for any values b_1, b_2, \dots, b_5 ? Completely fill in the circle next to the correct answer.

Concepts: This question checks whether you can determine consistency of a system of equations (e.g., by comparing dimension of the column space and dimension of \vec{b}).

Answer: \vec{b} will be a vector in \mathbb{R}^5 , but column-space of A has dimension at most 4...

- (e) (4 points) Assuming a solution exists for the new system of equations in part (d), will the solution be unique? Justify your answer by showing work to support your conclusion.

Concepts: This question checks whether you understand how nullspace relates to uniqueness of a solution to a consistent system of linear equations.

Answer: Yes, the solution will be unique.

PRINT your name and student ID: _____

7. Dynamical Systems (26 points)

Define matrices $Q, R \in \mathbb{R}^{2 \times 2}$ according to

$$Q = \begin{bmatrix} 0 & 3/4 \\ 1 & 1/4 \end{bmatrix}, \quad R = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

- (a) (5 points) Find the eigenvalues for the matrix Q .

Concepts: Do you know how to find the eigenvalues of a 2x2 matrix?

Answer: The eigenvalues are $\lambda_1 = 1, \lambda_2 = -3/4$.

- (b) (4 points) Consider a system with state vector $\vec{x}[n] \in \mathbb{R}^2$ at time $n \geq 1$ given by

$$\vec{x}[n] = Q\vec{x}[n-1].$$

Is there a non-zero vector \vec{x} satisfying $\vec{x} = Q\vec{x}$? If yes, give one such vector.


Concepts: Can you find an eigenvector corresponding to a given eigenvalue (1 in this case)?

Answer: Yes; $\vec{x} = [3/4, 1]^T$.

- (c) (3 points) Draw the state-transition diagram for the system in part (b). Label your nodes "A" and "B".

Concepts: Do you know how to draw a state-transition diagram for a system of linear equations?

Answer:



- (d) (4 points) Now, consider a system with state vector $\vec{w}[n] \in \mathbb{R}^2$ at time $n \geq 1$ given by:

$$\vec{w}[n] = \begin{cases} Q\vec{w}[n-1] & \text{if } n \text{ is odd} \\ R\vec{w}[n-1] & \text{if } n \text{ is even.} \end{cases}$$

Write expressions for $\vec{w}[1]$, $\vec{w}[2]$, $\vec{w}[3]$ and $\vec{w}[4]$ in terms of $\vec{w}[0]$ and Q and R . Write each answer in the form of a matrix-vector product.

Concepts: Given a description of a dynamical system, can you write out the state vectors at given time points in terms of the initial state and the transition matrices?

Answer:

$$\vec{w}[1] = Q\vec{w}[0], \quad \vec{w}[2] = RQ\vec{w}[0], \quad \vec{w}[3] = Q(RQ)\vec{w}[0], \quad \vec{w}[4] = (RQ)^2\vec{w}[0].$$

- (e) (10 points) Suppose we start the system of part (d) with state $\vec{w}[0] = [11/14 \quad 3/14]^T$. Find expressions for \vec{w}_{even} and \vec{w}_{odd} , which are defined according to

$$\vec{w}_{\text{even}} = \lim_{k \rightarrow \infty} \vec{w}[2k], \quad \vec{w}_{\text{odd}} = \lim_{k \rightarrow \infty} \vec{w}[2k+1].$$

In words, \vec{w}_{even} and \vec{w}_{odd} describe the long-term behavior of the system at even and odd time-instants, respectively. (Hint: you can avoid computation by thinking about the system at even time-instants in terms of a state-transition diagram.)

Concepts: Following the hint, you should consider the dynamical system $\vec{w}[2k] = (RQ)^k \vec{w}[0]$ for $k \geq 1$. This is just like the dynamical systems you have considered previously, with transition matrix (RQ) . If you compute this matrix product and draw the state diagram, you will find something that looks nearly identical to the page-rank example from lecture. So, this question tests whether you can recognize a familiar problem, perhaps presented in a slightly unfamiliar form (but guided by a hint).

Answer:

$$\vec{w}_{\text{even}} = \lim_{k \rightarrow \infty} \vec{w}[2k] = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad \text{and, } \vec{w}_{\text{odd}} = Q\vec{w}_{\text{even}} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

PRINT your name and student ID: _____

8. Linearly Independent Solutions (5 points)

Let $A \in \mathbb{R}^{17 \times 32}$ satisfy $\dim(C(A)) = 9$, where $C(A)$ denotes the column-space of A . How many linearly independent solutions can be found to the system of equations $A\vec{x} = \vec{0}$?

Note: Be careful. You are not being asked how many solutions exist for this system of equations, but rather how many *linearly independent solutions* can be found. You may just give a numerical answer; no work is required.

Concepts: Do you know (i) definition of dimension of a subspace (equal to max number of linearly independent vectors in a subspace); (ii) definition of null-space; and (iii) how dimension of null-space and column-space are related to matrix dimensions (i.e., rank nullity theorem)?

Answer: The number of linearly independent solutions is equal to the dimension of $N(A)$, which is the maximum number of linearly independent solutions to the equation $A\vec{x} = 0$, by definition. Hence, we use the rank-nullity theorem to compute.

PRINT your name and student ID: _____

9. Inverses and Transposes (8 points)

Given an invertible matrix $A \in \mathbb{R}^{n \times n}$, use the definition of matrix inverse to prove that

$$(A^T)^{-1} = (A^{-1})^T.$$

Concepts: Do you know the definition of matrix inverse (i.e., $AA^{-1} = A^{-1}A = I$)? Do you remember what happens when you take transpose of a matrix product?

PRINT your name and student ID: _____

10. Orthogonal Complements (16 points)

Consider the vector space \mathbb{R}^n , and let \mathbb{U} be a subspace of \mathbb{R}^n . We define the set $\mathbb{U}^\perp \subset \mathbb{R}^n$, called the *orthogonal complement* of \mathbb{U} , according to

$$\mathbb{U}^\perp = \{\vec{x} \in \mathbb{R}^n \mid \vec{u}^\top \vec{x} = 0 \text{ for all } \vec{u} \in \mathbb{U}\}.$$

- (a) (4 points) Show that \mathbb{U}^\perp is a subspace of \mathbb{R}^n .

Concepts: Do you know the definition of a subspace, and can you verify it on a given example?

Answer: We should show that \mathbb{U}^\perp is closed under scalar multiplication and vector addition.

- (b) (4 points) Find a concise expression for the intersection $\mathbb{U} \cap \mathbb{U}^\perp$. Justify your answer.

Concepts: You saw the operation \cap for subspaces in your homework; can you use definitions to compute it for a specific example?

Answer: If $\vec{x} \in \mathbb{U}^\perp$, then $\vec{x}^\top \vec{u} = 0$ for any choice of $\vec{u} \in \mathbb{U}$. In particular, if $\vec{x} \in \mathbb{U}^\perp \cap \mathbb{U}$, then we must have

$$0 = \vec{x}^\top \vec{x} = \sum_{i=1}^n x_i^2$$

Thus...

- (c) (6 points) Working in dimension $n = 3$, consider the subspace

$$\mathbb{U} = \text{span} \left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} \right\}.$$

Find a basis for \mathbb{U}^\perp .

Concepts: Can you formulate the problem of computing \mathbb{U}^\perp as a system of linear equations and solve? This is almost identical to how we compute the nullspace of a matrix: we formulate an appropriate system of linear equations, and then solve.

Answer: The vector $[-1, -1, 1]^T$ is a basis for \mathbb{U}^\perp .

- (d) (2 points) For the subspaces \mathbb{U} and \mathbb{U}^\perp of part (c), show that $\mathbb{U} + \mathbb{U}^\perp = \mathbb{R}^3$.

Concepts: Do you know that three linearly independent vectors in \mathbb{R}^3 will span \mathbb{R}^3 ?